

#### **Absolute Dermatology & Skin Cancer Center**

5207 Hickory Park Drive, Suite A, Glen Allen, VA 23059 7300 Ashlake Parkway, Suite #102, Chesterfield, VA 23832 T: (804) 326-4448 F: (309) 326-4947 www.absolutedermva.com

## BENIGN RESULTS WILL BE RELEASED TO THE PATIENT PORTAL. IF YOU HAVE NOT HEARD FROM US IN 2 WEEKS, PLEASE CHECK YOUR PORTAL FOR YOUR RESULTS.

#### **Minor Procedure Wound Care Instructions**

## Biopsy, Removal, or ED&C ("Burn & Scrape")

- Keep the wound as dry as possible for 24 hours. You then may begin bathing but avoid lakes, rivers, hot tubs or public swimming pools for at least one week. Avoid strenuous exercise for at least 24 hours.
- Clean the wound with soap and warm water once or twice daily. Apply a moderate amount of Vaseline or Polysporin (NOT NEOSPORIN) and a fresh bandage each time until healed.
- Some patients develop an irritant or allergic reaction to bandages (adhesive material). If this occurs, use paper tape and gauze.
- After you finish wearing the bandage it is helpful to rub in a tiny amount of Vaseline ointment twice daily for an additional week to keep the wound soft and free of germs.
- What to expect:
  - Effects of the numbing medication will wear off in 1-2 hours. If you need pain medication, Tylenol is the best choice as ibuprofen/aspirin can increase bleeding.
  - A small amount of bloody drainage is normal. If you experience active bleeding, remove the
    dressing and apply direct pressure for 10 minutes. If you continue to have active bleeding or
    develop pain, pus, or fever, please call the office.
  - Most healing wounds have a yellow appearance before the top layer of skin grows back and this is normal. Wounds on the lower leg and forearms heal very slowly and may take several weeks or more to completely heal.
  - o If pathology was sent, your results should be ready in 10-14 business days. If you have not received your results in 2 weeks, then you need to contact us.

# \_\_\_\_YOU HAVE A STITCH(ES) THAT NEED TO BE REMOVED IN\_\_\_\_DAYS AT OUR OFFICE/ AT HOME.

### Cryotherapy ("Freezing")

- It is normal for the site to be red, swollen and develop a blister the of your appointment. You may pop the blister with a sterile needle and drain the fluid if it becomes uncomfortable. Leave the roof of the blister in place.
- Cleanse the wound with gentle soap and water once a day. Do not scrub the area.
- Apply Vaseline or Polysporin (NOT NEOSPORIN) to the site twice daily. A bandage should be applied if the area becomes blistered or irritated.
- Some patients develop an irritant or allergic reaction to bandages. If this occurs, use paper tape and gauze.
- If you notice any signs of infection such as fever, foul smells, marked redness, swelling, or pain, call our office.

Please call us if you have any concerns or questions regarding your procedure or biopsy results!